




new

Pull out the toxins



As toxin exposure has been increasing at a rate faster than the human body is genetically equipped to manage, this foundational approach **improves patient response to common toxin exposure.**

Step 1 _ Open up detox pathways

Glutathione, known as the master antioxidant for its ability to recycle other antioxidants, is an essential component of the body's detox system. Glutathione plays an integral role in both Phase 1 and Phase 2 detoxification. Glutathione becomes depleted when the body is faced with an overwhelming amount of toxins.

The problem traditionally is that glutathione is poorly absorbed by the body in supplement form. **TriFortify®**, oral liposomal Glutathione, has overcome these barriers. Tri-Fortify® liposomal glutathione has been clinically researched and published in the European Journal of Clinical Nutrition in 2018.



Step 2 _ Remove the toxins

Mycotoxins

= Secondary mold metabolites released by spores in conditions when humidity and temperature are sufficient.



Glyphosate

= synthetic amino acid, most toxic herbicide in the USA. Glyphosate is classified as human carcinogen.

Heavy Metals

= Widespread contaminants like Cadmium, Mercury and Lead.



Step 3 _ Protect and replanish

Mineral status must be monitored during detoxification & chelation therapy.

Core Minerals™ provides restorage of minerals during chelation therapy with Mycopul™ or ToxinPul™



More information about Trifortify®, MycoPul™, ToxinPul™ and Core Minerals™ on the following pages.

Tri-Fortify Watermelon® or Orange®




indication	Detoxification with glutathione in high bioavailable formulation, powerful antioxidant, Natural Killer Cell support	
dosage	1 teaspoon (1 pack) per day, away from food	
packaging	236 ml per tube or 20 packs per box	
composition (amount per 1 teaspoon)	Glutathione Liposomal Vitamin C	450 mg 50 mg

All information is exclusively aimed at and released to an audience of health care professionals.


Glutathione levels

Increase in red blood cell levels (Erythrocytes)




Oxidative stress markers

Oxidized / Reduced GSH




Immune function

Natural Killer Cell activity



Lipid Peroxidation

(Reduced Cellular Membrane Oxidation)



Published research : Sinha, R., Sinha, I., Calcagnotto, A., Trushin, N. Oral supplementation with liposomal glutathione elevates body stores of glutathione and markers of immune function. Eur J Clin Nutr. 2018 Jan;72(1):105-111.

MycoPul™



indication	MycoPul is a combination of several binders, targeted to eliminate mycotoxins (toxins produced by molds), heavy metals and environmental toxins. MycoPul includes a patented, purified and natural form of zeolite.	
dosage	1 caps per day 1 hour before or 2 hours after eating	
packaging	30 caps per container	
composition (amount per 1 cap)	Activated Charcoal	150 mg
	Fulvic acid	100 mg
	Black humate powder	100 mg
	Zeolite Clinoptilolite (G-Pur®)	100 mg
	Microchitosan	100 mg
	Bamboo extract	50 mg

All information is exclusively aimed at and released to an audience of health care professionals.

MycoPul™

Main constituents



Humic/Fluvic powder blend with trace minerals	Mycotoxine binder, especially aflatoxines
Activated Charcoal	Removal of ochratoxines ¹ , aflatoxines, trichothecenes and zeralenone Activated Charcoal is an amorphous form of carbon prepared from incomplete combustion and activated by gas flow to create pores
G-Pur purified Zeolite Clinoptilolite (patented)^{2,3}	Removes aflatoxins, ammonia, histamine, and heavy metals Structure differs from other zeolite forms: <ul style="list-style-type: none"> • Certified purity • stable in acid pH environment • High ion exchange capacity • ↑ Silica/Alum ratio (4/1)
Microchitosan	Support of healthy removal of fungus from the body Removal of Ochratoxines and heavy metals ^{4,5} Microchitosan adsorbs metal ions through forming complexes ⁶ Antimicrobial activity Microchitosan binds to the negatively charged bacterial cell wall causing disruption of the cell, thus altering the membrane permeability
Silica (from bamboo extract)^{7,8,9,10,11}	

MycoPul™

References



- ¹ El Khoury R, Choque E, El Khoury A, et al. Ota prevention and detoxification by actinobacterial strains and activated carbon fibers: Preliminary results. *Toxins (Basel)*. 2018;10(4):1-16. doi:10.3390/toxins10040137
- ² Smyrna HP. G-PUR / Purified Clinoptilolite.; 2017.
- ³ Pavelić SK, Medica JS, Gumbarević D, Filošević A, Pržulj N, Pavelić K. Critical review on zeolite clinoptilolite safety and medical applications in vivo. *Front Pharmacol*. 2018;9(NOV):1-15.
- ⁴ Zhang L, Zeng Y, Cheng Z. Removal of heavy metal ions using chitosan and modified chitosan: A review. *J Mol Liq*. 2016;214:175-191.
- ⁵ Min LL, Zhong L Bin, Zheng YM, Liu Q, Yuan ZH, Yang LM. Functionalized chitosan electrospun nanofiber for effective removal of trace arsenate from water. *Sci Rep*. 2016;6(April).
- ⁶ Zia Q, Tabassum M, Gong H, Li J. A Review on Chitosan for the Removal of Heavy Metals Ions. *J Fiber Bioeng Informatics*. 2019;12(3):103-128.
- ⁷ Jacqmin-Gadda H, Commenges D, Letenneur L, Dartigues JF. Silica and aluminum in drinking water and cognitive impairment in the elderly. *Epidemiology*. 1996;7(3):281-285.
- ⁸ Rondeau V, Jacqmin-Gadda H, Commenges D, Helmer C, Dartigues JF. Aluminum and silica in drinking water and the risk of Alzheimer's disease or cognitive decline: Findings from 15-year follow-up of the PAQUID cohort. *Am J Epidemiol*. 2009;169(4):489-496.
- ⁹ Davenward S, Bentham P, Wright J, et al. Silicon-rich mineral water as a non-invasive test of the "aluminum hypothesis" in Alzheimer's disease. *J Alzheimer's Dis*. 2013;33(2):423-430.
- ¹⁰ Dalton AJ. Commentary on "Non-invasive therapy to reduce the body burden of aluminium in Alzheimer's disease." *J Alzheimer's Dis*. 2006;10(1):29-31.
- ¹¹ Jones K, Linhart C, Hawkins C, Exley C. Urinary Excretion of Aluminium and Silicon in Secondary Progressive Multiple Sclerosis. *EBioMedicine*. 2017;26:60-67.

ToxinPul™



indication	Elimination of heavy metals. Humic acid pulls heavy metals and toxins from the blood and elimination happens through the intestine.	
dosage	3 caps once daily, 1 hour before or 2 hours after food	
packaging	90 vegecaps per container	
composition (amount per 3 vegecaps)	Taurine	500 mg
	Dandelion root (Taraxacum officinale)	500 mg
	Fulvic acid	200 mg
	Humic acid	150 mg
	Cilantro leaf extract (Coriandrum sativum)	120 mg
	Vitamin C (as Ascorbic acid)	100 mg
	Quercetin (Quercefit™)	100 mg
	Chlorella powder (Chlorella vulgaris)	90 mg

All information is exclusively aimed at and released to an audience of health care professionals.

ToxinPul™

Main constituents



Humic Acid ^{1,2}	
Fulvic Acid ³	Targeted binders for multiple chemicals including pesticides (glyphosate), herbicides, and heavy metals
Silica	Silica to aid in binding aluminum and prevent accumulation of aluminum in the body)
Cilantro (Coriandrum Sativum) ^{4,5}	Cilantro to support mobilization of toxins in the body to increase removal
Chlorella (Chlorella Vulgaris) ⁶	Binding of toxins in both the GI tract and systemically
Dandelion (Taraxacum Officinale)	Dandelion for healthy liver support and protection from herbicides
Quercefit (highly absorbable quercetin) ^{7,8}	Quercefit™ (20x more absorbable form of Quercetin) to support healthy liver and kidney function
Taurine	Taurine to support bile flow for better toxin removal and a source of sulfur needed for detoxification pathways
Vitamin C ⁹	Vitamin C for antioxidant protection and protection from harm from mycotoxins

ToxinPul™

References



- ¹ Shehata AA, Kühnert M, Haufe S, Krüger M. Neutralization of the antimicrobial effect of glyphosate by humic acid in vitro. *Chemosphere*. 2014;104:258–261.
- ² Gondar D, Iglesias A, López R, Fiol S, Antelo JM, Arce F. Copper binding by peat fulvic and humic acids extracted from two horizons of an ombrotrophic peat bog. *Chemosphere*. 2006;63(1):82–88.
- ³ Glynn AW. Fulvic and humic acids decrease the absorption of cadmium in the rat intestine. *Arch Toxicol*. 1995;70(1):28–33.
- ⁴ Aga M, Iwaki K, Ushio S, et al. Preventive effect of *Coriandrum sativum* (Chinese parsley) on aluminum deposition in ICR mice. *Nat Med*. 2002;56(5):187–190.
- ⁵ Sharma V, Kansal L, Sharma A. Prophylactic efficacy of *coriandrum sativum* (coriander) on testis of lead-exposed mice. *Biol Trace Elem Res*. 2010;136(3):337–354.
- ⁶ Gerlach A. Oral Application of Charcoal and Humic acids to Dairy Cows Influences *Clostridium botulinum* Blood Serum Antibody Level and Glyphosate Excretion in Urine. *J Clin Toxicol*. 2014;04(02):1–8.
- ⁷ Soudani N, Chaâbane M, Ghorbel I, Elwej A, Boudawara T. Glyphosate disrupts redox status and up-regulates metallothionein I and II genes expression in the liver of adult rats. Alleviation by quercetin. *Gen Physiol Biophys*. 2014;31(4):375–382.
- ⁸ Riva A, Ronchi M, Petrangolini G, Bosio S, Allegrini P. Improved Oral Absorption of Quercetin from Quercetin Phytosome®, a New Delivery System Based on Food Grade Lecithin. *Eur J Drug Metab Pharmacokinet*. 2019;44(2):169–177.
- ⁹ Gehin A, Guyon C, Nicod L. Glyphosate-induced antioxidant imbalance in HaCaT: The protective effect of Vitamins C and E. *Environ Toxicol Pharmacol*. 2006;22(1):27–34.

Core Minerals™



indication	Balanced mineral complex without iron: restorage of minerals during chelation therapy	
dosage	1 x 4 caps per day, 12 hours after chelation	
packaging	120 vegecaps per container	
composition (amount per 4 vegecaps)	Calcium (as Calcium citrate + malate)	300 mg
	Magnesium (as Magnesium glycinate)	300 mg
	Potassium (as Potassium citrate)	99 mg
	Zinc (as Zinc picolinate)	15 mg
	Copper (as Copper glycinate)	2 mg
	Manganese (as Manganese glycinate)	2 mg
	Boron (as Boron picolinate)	2 mg
	Chromium (as Chromium polynicotinate)	200 µg
	Iodine	150 µg
	Selenium (as Selenium citrate)	100 µg
	Molybdenum (as Molybdenum picolinate)	100 µg
	Vanadium (as Vanadylsulfate)	100 µg
	Vitamin D3 (as Cholecalciferol)	5 µg

All information is exclusively aimed at and released to an audience of health care professionals.