



focus


Doctor's Biome: Protect your gate-keepers the way they protect you



Within the context of optimal health, the importance of our gut should not be underestimated. The scientific team of Doctor's Biome conducted after various methods of research (see p. 4-11) the ideal probiotic supplement.

Starting from this belief, the company was very mindful of its product's ingredients throughout the process and the choice of probiotics is based on recent scientific studies. They thus also concluded it is important for the product to be **100% organic**.

What makes the product perhaps the most progressive is yet something else. The probiotics are physiologically active during the time of manufacture and storage. What does this mean? While the majority of probiotic companies will not tell you if the CFU (colony forming units) of their products will all survive rehydration, we guarantee that **the CFUs of actually living probiotics grow and thus proliferate**. Their **27 billion** CFUs at the time of production is typically **50-100 billion** CFUs when it reaches the consumer's hands.



So, the composition of their product ensures that the living bacteria keep increasing in number, meaning that Doctor's Biome becomes more and more efficient by time. It took almost **3 years and 3 nutritional scientists**, but they managed to create a fully organic product that obtains the desired effects of a healthy gut system without disturbing the fragile and important balance of the gut biome. Simply because we, Nutrined as well as the scientific team of Doctor's Biome, believe that **good health starts from a good gut**.

Your body sends you a signal

In some cases, our gut system is no longer optimally efficient due to age or a weakened immune system – perhaps caused by regular intake of antibiotics. In this case, one can put a lot of effort in life-style and diet, if the gut is unable to do its job as its supposed to, one will not obtain the desired results. Unpleasant problems such as diarrhea, flatulence, constipation or bloating might be experienced in this case. Such problems, ranging from occasional to alarming frequency, is your body sending you a signal. Quick solutions may include keeping track of your diet with a food diary, avoiding certain meals or taking dietary supplements.

1. Infections

If there is an active infection a supplement is required. In the case of Clostridium Difficile (C. difficile), for example. This bacterium can cause a life-threatening, uncontrolled diarrhea. Our good-natured bacteria are then no longer strong enough to defend themselves against the bad-natured ones. In many cases, C. difficile is caused by an adverse effect of taking antibiotics. (Read the research on p. 4-11)

2. Optimal health

However, not only in the case of an infection Doctor's Biome can be the solution. In a proactive way on the contrary, when one aims for optimal health, the product can make the difference. When we think of the chief players that maintain our bodily system, we easily consider the heart or the brain the ones taking charge. However, without dishonoring these major stakeholders, we often tend to forget a part of our body, one that is arguably at the top of our physical ladder: **our gut**.


The body being much smarter than we would superficially suspect it to be is nothing new. Apart from physical wounds that it manages to heal time after time or mental toughness that we rarely run out of, is there **an entire system operating behind the scenes** in which each organ has its own role.

Let us start at the beginning: **food**. One eats healthy as we've been taught it is good for us. But what happens to food the moment it enters our body?

Guarding the fort

The **digestion process** starts at the oesophagus, then down to the stomach, then the small intestine and later to the large intestine.

The **small intestine** is an important part of the process: here, the food is, literally, kneaded through the muscle layer for about 4 to 8 hours. Digestive enzymes are added to the food and usable fats, minerals, sugars and proteins are filtered out. Those nutrients are necessary to build or repair the body but some also operate as all-around fuel. It is the liver's job to send all the nutrients to the right location in the body.



Once the small intestine's job is done, the food moves on to the **large intestine** where the good bacteria gather. Think of it as a balanced-out pool, also known as our microbiome, in our body, where good-natured bacteria guard the fort. Lastly there is the final step: the stool.



Make a pact with your gut

Now, why is it thus crucial to understand the role of your gut if one wants to obtain optimal health? You could think of the gut as our body's gate-keepers. Not only do they digest and filter our food, it is also where toxins and harmful bacteria enter our body. **70% of our immune system is located in the gut.**

In addition, due to climate change and increasing pesticide use, our food is not as nutritious as it used to be. For this reason, it is even more important that our gut manages to filter out the beneficial substances.

Perhaps you had not yet considered this as one of your urgent to do's, but it is very crucial to be kind to your gut within the framework of optimal health. Exercise, (as this literally puts your system in motion), eat as much fibre as possible (fruit, vegetables and nuts are an accessible way to do so), drink water and avoid stress. But sometimes our gut system is disrupted by factors beyond our control, and that's when **we need an extra push.**

**70% of
our immune
system is
located in
the gut.**





Original Article

DOI: 10.36959/621/560

Complete Inhibition of *Clostridium difficile* with a Probiotic Juice Beverage

Howard F Robins^{1*} and A Reza Kamarei²¹Chief Medical Officer, Doctor's Biome, Hauppauge, NY, USA²Chief Science & Technology Officer, Doctor's Biome, Hauppauge, NY, USA

Abstract

Background & Aims: *Clostridium difficile* (“*C. difficile*” or “*C. diff*”) is a Gram-positive, spore-forming, anaerobic, motile bacterium which produces two types of toxins, enterotoxin A and cytotoxin B. *Clostridium difficile* Infection (CDI) causes life-threatening diarrhea which is usually a side-effect of taking antibiotics. *C. diff* can easily spread from person to person and is considered a major health threat. In 2017, there were an estimated 223,900 cases in hospitalized patients and 12,800 deaths in the United States. There has been considerable scientific research in using various probiotic strains against *C. diff* with diverse observations and inconsistent results. Our first goal was to develop an “optimum probiotic product” to inhibit growth and pathogenicity of *C. diff*. Our second goal was to prove an inhibitory effect of bioactive compounds (secreted by the 15 strains of probiotic bacteria into the juice) on *C. diff*.

Methods: For optimum profile of probiotics, we considered positive results from specific strains in the published literature and commercially-available probiotic bacteria from credible global suppliers. As a result, we chose a blend of five strains of Bifidobacteria and ten strains of Lactobacilli. For an optimum carrier, we chose a proprietary blend of organic green vegetable and fruit juices. Upon culture propagation and strain verification of *Clostridium difficile* ATCC #9689, inoculum was prepared using reinforced clostridial broth and a sample was prepared using centrifuged, microfiltered supernatant of Doctor's Biome Colon Health (DBCH). The inoculated *control* was serially diluted with sterile 0.1% Peptone Buffer. The inoculated *sample* was serially diluted with the centrifuged, filtered supernatant of DBCH. The serially diluted control and diluted sample were transferred to agar plates and incubated anaerobically at 35-37 °C for 18-24 hours.

Results: Serially diluted agar plates of the control showed presence of *C. diff* colony forming units, which were biochemically identified via VITEK ANC. Serially diluted agar plates of the sample did not show any *C. diff* colony forming units due to the inhibitory effect of the DBCH bioactive compounds.

Conclusions: We believe inhibition of *C. diff* by DBCH probiotics should be considered from both a quantitative and qualitative points of view. From a quantitative point of view, a general mechanism of microbial inhibition is through the “competitive exclusion principle”. This means that when two species compete in a limited environment (e.g., colon) and compete for the same limited amount of nutrients, the species that has advantage over the others (e.g., larger numbers) will dominate the environment and lead to the exclusion of the weaker competitor. From a qualitative point of view, it seems that the chosen blend of Bifidobacteria and Lactobacilli have released some bioactive compounds into the juice that has completely inhibited growth of *C. diff*. We conclude that the bioactive compounds secreted from the 15 probiotic strains into the organic vegetable and fruit juice has played a decisive inhibitory role against growth of *C. diff*.

Keywords

Clostridium difficile, *C. difficile*, *C. diff*., *Clostridium difficile* infection (CDI), Probiotics, Multi-strain probiotics, Liquid probiotics

Introduction

Clostridium difficile, also known as “*C. difficile*” or “*C. diff*” is a Gram-positive, spore-forming, anaerobic, motile bacterium (Figure 1). *Clostridium difficile* cells show optimum growth on blood agar at human body temperatures in the absence of oxygen. *Clostridium difficile* produces two types of toxins, enterotoxin A and cytotoxin B.

C. difficile is a bacterium that causes life-threatening diarrhea. It is usually caused as an adverse effect of taking

***Corresponding author:** Dr. Howard F Robins, Chief Medical Officer, Doctor's Biome (Newgen 27 LLC), 80 Davids Dr., Unit 2, Hauppauge, NY 11788, USA, Tel: (631)-240-3066, Cell: (516)-967-1009

Accepted: August 06, 2020

Published online: August 08, 2020

Citation: Robins HF, Kamarei AR (2020) Complete Inhibition of *Clostridium difficile* with a Probiotic Juice Beverage. J Gastroenterol Res 4(1):104-111



Figure 1: *Clostridium difficile* (“*C. diff.*”).

Source: US center for disease control and prevention (CDC).

antibiotics. It is considered a major health threat and listed as an “Urgent Threat” by the Center for Disease Control and Prevention (CDC) [1]. According to the CDC symptoms may begin within a few days or several weeks after beginning or following the taking of antibiotics or when people touch surfaces that are contaminated with feces from an infected person. Symptoms of *Clostridium difficile* Infection (CDI) include diarrhea, loose or watery stools lasting for several days, fever, stomach tenderness, loss of appetite and nausea. Frequent recurrent and duplicate episodes are common with one in six patients with CDI occurring again within two to eight weeks [1].

These infections are common in people 65 and older (80+%) and in children who take antibiotics and receive outpatient medical care or people who stay in hospitals and nursing homes for a long period of time [1,2]. Also, it is common in patients with weakened immune systems on antibiotics or with previous infection with *C. diff.* One out of eleven patients diagnosed with CDI will die within a month [1].

The estimated national burden of *C. difficile* infection (CDI) was 476,400 cases in 2011 and 462,100 in 2017 with an average number of deaths around 28,000 per year [3].

CDI puts a tremendous strain on the healthcare system. The total financial burden of CDI inpatient management was estimated to be \$5.4-6.3 billion USD per year. This includes 2.4 million days of hospital stays and growing in number yearly [3-6].

With the increased use of appropriate and inappropriate prescriptions of antibiotics both in hospital and outpatient care the financial burden from CDI has risen over the past decade [4,7-9].

The scientific definition of the term ‘probiotic’ was proposed in a 2001 joint report of the Food and Agriculture Organization of the United Nations and the World Health Organization (FAO/WHO), and confirmed by an expert panel con-

vened by International Scientific Association for Probiotics and Prebiotics (ISAPP) held in 2013. The currently accepted definition of probiotics is: “Live microorganisms that, when administered in adequate amounts, confer a health benefit on the host”. This means if the microorganisms are not “live”, they cannot be considered probiotic and if their amount is not adequate enough to confer a health benefit, they cannot be considered probiotic either.

A review of the literature shows it is replete with hundreds of published papers showing various positive and some mixed results on the use of specific probiotic strains to help prevent and help treat CDI [10-15].

The probiotics used in most of the consumer available preparations are usually taken from a limited range of organisms. As this area of study has become more important and prevalent, new strains have come under investigation.

“Next Generation Probiotics” (NGP) are bacteria that due to their development act as live biotherapeutic products and lend themselves to act more like pharmaceuticals than traditional food supplements [16,17].

Despite the above findings reported in the literature, McDonald, *et al.*, in 2017 *update on Clinical Practice Guidelines for Clostridium difficile Infection in Adults and Children*, using earlier publications up to 2013 and referring to issues such as differences in probiotic formulations studied, duration of probiotic administration, definitions of CDI, duration of study follow-up, inclusion of patients not typically considered at high risk for CDI, and the potential for organisms in probiotic formulations which can cause infections in hospitalized patients, concluded that there is insufficient data at this time to recommend administration of probiotics for primary prevention of CDI outside of clinical trials of probiotics for primary prevention of CDI. Yet the literature is replete with numerous studies showing that specific strains of patented bacteria have had beneficial results in the prevention and treatment of CDI.

The Harvard School of Public Health says “Because probiotics fall under the category of supplements and not food, they are not regulated by the Food and Drug Administration in the U.S. This means that unless the supplement company voluntarily discloses information on quality, such as carrying the USP seal that provides standards for quality and purity, a probiotic pill may not contain the amounts listed on the label or even guarantee that the bacteria are alive and active at the time of use” [18].

Several studies have been performed in an attempt to find a way to improve survival of probiotics before reaching the large intestine with limited and minimal success. This is one of the main concerns over capsule, powder, tablet and spore probiotics. Less than 59% survival using unique and somewhat expensive measures have been employed, though rarely found and used in most probiotic preparations currently available [19-24].

The FDA states “Many dietary supplements, often marketed as “probiotics,” include claims that the live microbial ingredient, formulated in adequate amounts, will confer a health benefit on consumers. The weight of microbial dietary ingredient in a product represents the product’s total cellular mass, consisting of both live and dead microorganisms, and therefore does not necessarily correlate with the number of viable microorganisms in that product” [25].

Aims of the Study

It became evident that certain strains of NGP “Smart Bacteria” (SB) achieved consistently the best results in helping to prevent and assist in CDI amelioration along with the usual treatment regimens. SB can sense they’re attached to our intestinal cells, and then remodel their expression of specific genes, including those involved in metabolism, to beneficially exploit our cells and colonize our gut [26,27].

Bacteria have evolved to develop remarkable systems that can sense neighboring cells called “target” cells, and can initiate upon contact a series of changes that enhance their ability to survive and grow at the expense of these “target” cells. Smart Bacteria do this by employing a mechanism called “Horizontal Gene Transfer” (HGT). This is a very interesting and complex mechanism. Simply put, HGT shares genetic material between organisms that are not in a parent-offspring relationship and is a widely recognized mechanism for adaptation in bacteria [28-33].

With this concern in mind, NGP SB’s were created to survive the stomach acids and digestive enzymes 80% or more [34]. Also, to ensure that they are “living” and thriving at the time of ingestion an appropriate environment and media, a 100% organic vegetable-fruit juice needed to be found meeting all these needs including the range of acidity needed for survival and growth. The NGP SB’s that were chosen needed to be compatible with each other so as not to overwhelm each other’s growth and benefits. They also needed to be in a sufficient number of Colony Forming Units (CFU) to repopulate the colon and attack the CDI effectively.

Considering diverse observations and inconsistent results of various probiotic strains on *Clostridium difficile*, it was not



Figure 2: Doctor’s biome colon health™.

(A patent-pending blend of 15 strains of probiotics in organic vegetable/fruit juice).

possible to choose one single, most effective strain for inhibition of *C. difficile*. Therefore, we decided to develop an “optimum probiotic product” to inhibit growth and pathogenicity of *C. difficile*. We named it *Doctor’s Biome Colon Health™* or DBCH (Figure 2). By definition, DBCH should have the optimum profile of probiotic bacteria and be an optimum carrier for the chosen probiotic bacteria. The immediate goal of our study was to prove the inhibitory effect of DBCH bioactive compounds (secreted by the 15 strains of NG Smart probiotic bacteria in the juice) on *C. difficile*.

Materials and Methods

Optimum profile of probiotics

Bifidobacteria and Lactobacilli are broadly recognized for their key roles in the human intestinal microflora throughout life. A high proportion of bifidobacteria and lactobacilli in the intestinal tract is considered beneficial to health. Considering the results of published literature mentioned above and reviewing commercially-available probiotic bacteria from credible global suppliers, we chose a blend of five strains of Bifidobacteria (Table 1a) and ten strains of Lactobacilli (Table 1b) from DuPont Nutrition & Biosciences.

Optimum carrier

In search of an optimum carrier (also known as excipients or medium), we chose a proprietary blend of organic green vegetable and fruit juices consisting of 100% organic diluted mint juice, cucumber juice, apple juice, lettuce juice, kale juice, celery juice and lemon juice. Sensory evaluation of DBCH (blend of probiotics in blend of organic juices) showed a pleasant taste of this product to consumers across the board.

Table 1a: Bifidobacteria of doctor's biome colon health.

(registered trademarks of DuPont-Danisco)

Bifidobacterium bifidum Bb-06	This is a specially selected strains of Bifidobacterium bifidum that could be used with a variety of excipients.
Bifidobacterium breve Bb-03	This strain is a specially selected strain of Bifidobacterium breve. B. breve (Bb-03) has demonstrated very good adhesion to human epithelial cell lines (HT-29) applied in <i>in vitro</i> studies. Bb-03 is tolerant to low pH conditions and survives the presence of bile at concentrations present in the duodenum.
Bifidobacterium lactis Bi-07*	This strain is a specially selected strain of Bifidobacterium lactis that could be part of a multiple strain custom blend with a variety of excipients. BI-07 has strong adhesion to intestinal cell lines.
Bifidobacterium longum BI-05	This is a specially selected strain of Bifidobacterium longum. <i>In vitro</i> studies have shown that B. longum BI-05 is tolerant to low pH conditions and survives the presence of bile at the concentrations present in the duodenum. B. longum BI-05 has demonstrated very good adhesion to human epithelial cell lines applied in <i>in-vitro</i> studies.
Bifidobacterium infantis Bi-26	This is a specially selected strain of Bifidobacterium longum subsp. infantis that could be part of a multiple strain custom blend with a variety of excipients.

Table 1b: Lactobacilli of doctor's biome colon health.

(registered trademarks of DuPont-Danisco)

Lactobacillus acidophilus La-14*	This strain is a specially selected strain of Lb. acidophilus. Well-suited for intestinal survival it has a high tolerance to gastrointestinal conditions (acid, bile, pepsin and pancreatin). LA-14 also has strong adhesion to intestinal cell lines. LA-14 may influence immune regulation as demonstrated by the induction of IL-12 and moderate induction of tumor necrosis factors <i>in vitro</i> .
Lactobacillus brevis Lbr-35	This is a specially selected strain of Lb. brevis that could be part of a multiple strain custom blend with a variety of excipients.
Lactobacillus bulgaricus Lb-87	This strain is a specially selected strain of Lb. bulgaricus. L. bulgaricus metabolizes and produces D (-)-lactic acid from four sugars: Lactose, glucose, fructose and mannose. LB-87 has the ability to rapidly convert lactose into lactic acid. LB-87 also has the ability to convert lactose into lactate, aiding in lactose digestion.
Lactobacillus casei Lc-11*	This strain is a specially selected strain of Lb. casei. LC-11 is well suited for intestinal survival and functionality. It is highly tolerant of acid and bile. LC-11 adheres strongly to intestinal Caco-2 cell lines and it inhibits common pathogens.
Lactobacillus gasseri Lg-36	This strain is a specially selected strain of Lb. gasseri that could be part of a multiple strain custom blend with a variety of excipients.
Lactobacillus paracasei Lpc-37*	This strain is a specially selected strain of Lb. paracasei. Lpc-37 displayed <i>in vitro</i> inhibition of selected pathogens including Salmonella typhimurium, Staphylococcus aureus, Escherichia coli and Listeria monocytogenes. A five-strain formulation including L. paracasei (Lpc-37) was found to maintain and more rapidly restore microbiota after antibiotic treatment. LPC-37 is well-suited for intestinal survival, it has a high tolerance to gastrointestinal conditions (acid and bile).
Lactobacillus plantarum Lp-115*	This strain is a specially selected strain of Lb. plantarum. <i>In vitro</i> studies have shown that L. plantarum (Lp-115) is extremely resistant to low pH conditions and survives the presence of bile at concentrations present in the duodenum. Lp-115 displayed <i>in vitro</i> inhibition of selected pathogens including Salmonella typhimurium, Staphylococcus aureus, Escherichia coli and Listeria monocytogenes. LP-115 is well-suited for intestinal survival; it has a high tolerance to gastrointestinal conditions (acid, bile, pepsin and pancreatin) and has strong adhesion to intestinal cell lines. LP-115 may improve specific immune response, as demonstrated in a human clinical study.
Lactobacillus reuteri 1E1	This is a specially selected strain of Lactobacillus reuteri that could be part of a multiple strain custom blend with a variety of excipients.
Lactobacillus rhamnosus Lr-32*	This strain is a specially selected strain of Lb. rhamnosus. L. rhamnosus (Lr-32) has shown anti-inflammatory properties, as demonstrated through significant protection against TNBS induced colitis in an animal model. L. rhamnosus (Lr-32) may influence immune regulation, as demonstrated by the increased induction of IL-10 <i>in vitro</i> .
Lactobacillus salivarius Ls-33*	This is a specially selected strain of Lb. salivarius. LS-33 is well-suited for intestinal survival, it has a high tolerance to gastrointestinal conditions (acid, bile, pepsin and pancreatin). LS-33 has a strong adhesion to intestinal cell lines, and ability to inhibit common pathogens. It also has a beneficial modulation of immune functions, it may influence immune regulation, as demonstrated by increased induction of IL-10 <i>in vitro</i> .

Furthermore, every batch of Doctor's Biome Colon Health is tested by an independent, FDA-registered accredited microbiology lab to confirm absence of pathogenic microorganisms.

Design of the Study

Culture propagation and strain verification

A sterile inoculating loop of *Clostridium difficile* ATCC #9689 was streaked onto a Reinforced Clostridial agar slant, broth tubes, and two Sheep Blood agar plates. The slant and broth tubes and one Sheep Blood agar plate were incubated anaerobically at 36-38 °C for 48 ± 2 hours. The second Sheep Blood agar plate was incubated aerobically at 36-38 °C for 48 ± 2 hours and served as the contamination check. Following incubation, the agar plates and tube were observed for colony growth and turbidity. The Sheep Blood agar plate incubated anaerobically was biochemically identified using the VITEK ANC card to verify purity.

Inoculum preparation

A well isolated colony from the Sheep Blood agar plate was used to inoculate a Reinforced Clostridial broth, which was then incubated anaerobically at 36-38 °C for 18 to 20 hours. Following incubation, the culture was stored overnight at 3 ± 1 °C.

Sample preparation

Six, individually packaged, intact 2-ounce DBCH glasses were composited into a sterile 500 mL glass bottle under a laminar flow hood. This bottle was mixed by gentle swirling and stored at ambient temperature (20-25 °C) until testing. At that time, the composited product was homogenized. Then 10 mL was pipetted into a test tube for a before (Figure 3) and after comparison photo with the centrifuged product. Follow-

ing centrifugation at 4500 RPM for 15 minutes, the supernatant liquid was filtered through a 0.22 µm filter to ensure that the supernatant liquid was free of cells. A 10 mL portion was pipetted into a test tube and photographed (Figure 4). The filtered sample was used as the test product **sample** and the 9 mL diluent tubes for its ten-fold serial dilution. The inoculated **control** was analyzed with sterile 0.1% Peptone Buffer.

Test procedure

Test procedure was performed according to the Compendium of Methods for the Microbiological Examination of Foods, 5th Edition, Chapter 33, Section 33.7 (Editors: Yvonne Salfinger and Mary Lou Tortorello, American Public Health Association).

Control test procedure: The inoculum used for the sample test procedure was vortexed for 5 seconds. 1 mL of the inoculum was transferred to a 9 mL tube of 0.1% Peptone Buffer and vortexed for 5 seconds. This served as the 10⁻¹ dilution tube. 1 mL from the 10⁻¹ dilution tube was transferred in duplicate into sterile Petri Dishes. With the same pipet, 1 mL was transferred into a second 9 mL tube containing the 0.1% Peptone Buffer for the 10⁻² dilution tube and vortexed for 5 seconds. 1 mL from the 10⁻² dilution tube was transferred in duplicate into sterile Petri Dishes. With the same pipet, 1 mL was transferred into a third 9 mL tube containing the 0.1% Peptone Buffer for the 10⁻³ dilution tube and vortexed for 5 seconds. Serial dilution was repeated until a 10⁻⁸ dilution plate was reached. Approximately 20 mL of tempered (44-46 °C) Reinforced Clostridial agar was added to the Petri Dishes and gently swirled to evenly disperse inoculum throughout the agar. After solidification, 15 mL of Reinforced Clostridial agar was overlaid onto each plate. The plates were incubated anaerobically in an upright position at 35-37 °C for 18-24




Figure 3: Doctor's biome colon health pre-centrifugation.



Figure 4: Doctor's biome colon health post-centrifugation/microfiltration.



Figure 5: Presence of *C. diff.* colony forming units when Peptone Buffer is used for serial dilution.

hours.

Sample test procedure: The refrigerated anaerobic chamber containing the inoculum was equilibrated to ambient temperature prior to inoculation. The inoculum was vortexed for 5 seconds to homogenize. 1 mL of the inoculum was transferred to a 9 mL tube of the supernatant test sample and vortexed for 5 seconds. This served as the 10^{-1} dilution tube. 1 mL from the 10^{-1} dilution tube was transferred in duplicate into sterile Petri Dishes. With the same pipet, 1 mL was transferred into a second 9 mL tube containing the supernatant test sample for the 10^{-2} dilution tube and vortexed for 5 seconds. 1 mL from the 10^{-2} dilution tube was transferred in duplicate into sterile Petri Dishes. With the same pipet, 1 mL will be transferred into a third 9 mL tube containing the supernatant test sample for the 10^{-3} dilution tube and vortexed for 5 seconds. Serial dilution was repeated until a 10^{-8} dilution plate was reached. Approximately 20 mL of tempered (44-46 °C) Reinforced Clostridial agar was added to the Petri Dishes and gently swirled to evenly disperse inoculums throughout the agar. After solidification, 15 mL of Reinforced Clostridial agar was overlaid onto each plate. The plates were incubated anaerobically in an upright position at 35-37 °C for 18-24 hours.

Results

Following incubation, photographs were taken of the se-



Figure 6: Absence of *C. diff.* colony forming units when doctor's biome colon health supernatant is used for serial dilution.

Table 2: Colony Counts of *Clostridium difficile*.

	Control		Sample	
Dilution	Rep 1	Rep 2	Rep 1	Rep 2
10^{-1}	TNTC	TNTC	0	0
10^{-2}	TNTC	TNTC	0	0
10^{-3}	TNTC	TNTC	0	0
10^{-4}	TNTC	TNTC	0	0
10^{-5}	TNTC	TNTC	0	0
10^{-6}	62	40	0	0
10^{-7}	3	1	0	0
10^{-8}	0	0	0	0

TNTC: Too numerous to count.

rially diluted agar plates of the **control** to visually show presence of *Clostridium difficile* colony forming units (Figure 5). The plates which exhibited less than 300 colonies were enumerated and representative colonies for *Clostridium difficile* were biochemically identified via VITEK ANC (Table 2).

Similarly, following incubation, photographs were taken of the serially diluted agar plates of the **sample** to visually show absence of *Clostridium difficile* colony forming units due to the inhibitory effect of the DBCH bioactive compounds (Figure 6 and Table 2).

Discussion and Conclusion

The obtained results of our *in-vitro* study are consistent with several published investigations in PubMed. We believe inhibition of *C. difficile* by DBCH probiotics should be considered from both quantitative and qualitative points of view:

From a quantitative point of view, a general mechanism of microbial inhibition is through the “competitive exclusion principle”. This means that when two species compete in a limited environment (e.g., colon) and compete for the same limited amount of nutrients, the species that has advantage over the others (e.g., larger numbers) will dominate the environment and lead to the exclusion of the weaker competitor. For example, in our study the probiotic dose of 27 billion live and active Colony Forming Units (CFU) for a daily serving of DBCH is considered among the higher quantities commercially available.

From a qualitative point of view, it seems that the chosen blend of *Bifidobacteria* and *Lactobacilli* have functioned in a complementary, additive and possibly synergistic way to completely inhibit growth of *C. difficile*. In other words, the chosen probiotics have released some bioactive compounds into the juice that has inhibited growth of *C. difficile*. One such compound could possibly be lactic acid (and possibly other organic acids) which can reduce the pH of the environment to an acidic range unfavorable for spore germination and/or growth of *C. difficile*. We can conclude that the bioactive compounds secreted from the chosen 15 probiotic strains into the organic vegetable and fruit juice has played a decisive inhibitory role against growth *C. difficile*.

The combined quantitative and qualitative properties of juice-based DBCH is expected to positively and impressively impact prevention and/or treatment of CDI.

Based on these results, and in line with the recent guidelines of American Gastroenterological Association on the role of probiotics in the management of gastrointestinal disorders [35], a prospective, randomized, double-blind clinical trial (based on Good Clinical Practice) on the safety and efficacy of DBCH for CDI patients is warranted.

Acknowledgment

Authors acknowledge the valuable contribution of Mary Lee Kluger and Andrew Liles from the independent Certified Laboratories (Melville, New York) for microbiological tests.

Financial Disclosure

Dr. Kamarei was paid as a consultant. Certified Laboratories was monetarily compensated by Doctor's Biome microbiological tests.

Conflict of Interest

Dr. Robins and Dr. Kamarei are partners in Doctor's Biome Company (Newgen 27 LLC).

References

1. Center for Disease Control: Healthcare associated diseases-community interface (HAIC).
2. Li N, Zeng B, Cai HE, et al. (2018) Cost-effective analysis of oral

antibiotics for the prevention of *clostridium difficile*-associated diarrhea in children and adolescents. *J Hosp Infect* 99: 469-474.

3. Guh AY, Mu Y, Winston LG, et al. (2020) Trends in U.S. burden of *clostridioides difficile* infection and outcomes. *N Engl J Med* 382: 1320-1330.
4. Dongmu Zhang, Vimalanand S Prabhu, Stephen W Marcella (2018) Attributable healthcare resource utilization and costs for patients with primary and recurrent *clostridium difficile* infection in the United States. *Clinical Infectious Disease* 66: 1326-1332.
5. Bond SE, Boutlis CS, Yeo WW, et al. (2017) The burden of healthcare-associated *clostridium difficile* infection in a non-metropolitan setting. *J Hosp Infect* 95: 387-393.
6. Heimann SM, Cruz Aguilar MR, Mellinshof S, et al. (2018) Economic burden and cost-effective management of *Clostridium difficile* infections. *Med Mal Infect* 48: 23-29.
7. Mundkur ML, Franklin J, Huybrechts KF, et al. (2018) Changes in outpatient use of antibiotics by adults in the United States 2006-2015. *Drug Saf* 41: 1333-1342.
8. Fleming-Dutra KE, Hersh AL, Shapiro DJ, et al. (2016) Prevalence of inappropriate antibiotic prescriptions among US ambulatory care visits, 2010-2011. *JAMA* 315: 1864-1873.
9. Donskey CJ (2017) *Clostridium difficile* in older adults. *Infect Dis Clin North Am* 31: 724-756.
10. Thad Wilkins, Jacqueline Sequoia (2017) Probiotics for gastrointestinal conditions: A summary of the evidence. *Am Family Physician* 96: 170-178.
11. Joshua Z Goldenberg, Christina Yap, Lyubov Lytvyn, et al. (2017) Probiotics for the prevention of *clostridium difficile*-associated diarrhea in adults and children. *Cochrane Database Sys Rev* 19.
12. John P Mills, Krishna Rao, Vincent B Young (2018) Probiotics for prevention of *clostridium difficile* infection. *Curr Opin Gastroenterol* 34: 3-10.
13. Johnston BC, Ma SS, Goldenberg JZ, et al. (2012) Probiotics for the prevention of *clostridium difficile*-associated diarrhea: A systematic review and meta-analysis. *Ann Intern Med* 157: 878-888.
14. Goldenberg JZ, Ma SS, Saxton JD, et al. (2013) Probiotics for the prevention of *clostridium difficile*-associated diarrhea in adults and children. *Cochrane Database Syst Rev* 31.
15. Goldstein EJC, Johnson SJ, Maziade PJ, et al. (2017) Probiotics and prevention of *clostridium difficile* infection. *Anaerobe* 45: 114-119.
16. O'Toole PW, Marchesi JR, Hill C (2017) Next-generation probiotics: The spectrum from probiotics to live biotherapeutics. *Nat Microbiol* 2: 17057.
17. Chang CJ, Lin TL, Tsai YL, et al. (2019) Next generation probiotics in disease amelioration. *J Food Drug Anal* 27: 615-622.
18. (2020) The microbiome. The nutrition source, Harvard School of Public Health.
19. Keller D, Verbruggen S, Cash H, et al. (2019) Spores of *bacillus coagulans* GBI-30, 6086 show high germination, survival and enzyme activity in a dynamic, computer-controlled in vitro model of the gastrointestinal tract. *Benef Microbes* 10: 77-87.
20. Maathuis AJ, Keller D, Farmer S (2010) Survival and metabolic activity of the GadenBC30 strain of *bacillus coagulans* in a dynamic in vitro model of the stomach and small intestine. *Benef Microbes* 1: 31-36.

21. Varankovich N, Martinez MF, Nickerson MT, et al. (2017) Survival of probiotics in pea protein-alginate microcapsules with or without chitosan coating during storage and in a simulated gastrointestinal environment. *Food Sci Biotechnol* 26: 189-194.
22. Rui Ji, Wu J, Zhang J, et al. (2019) Extending viability of bifidobacterium longum in chitosan-coated alginate microcapsules using emulsification and internal gelation encapsulation technology. *Front Microbiol* 10: 1389.
23. Ding WK, Shah NP (2009) An improved method of microencapsulation of probiotic bacteria for their stability in acidic and bile conditions during storage. *J Food Sci* 74: 53-61.
24. Ramos PE, Cerqueira MA, Teixeira JA, et al. (2018) Physiological protection of probiotic microcapsules by coatings. *Crit Rev Food Sci Nutr* 58: 1864-1877.
25. FDA policy regarding quantitative labeling of dietary supplements containing live microbials: Guidance for industry 21 CFR 10.115(g)(5)).
26. Landry BP, Tabor JJ (2017) Engineering diagnostic and therapeutic gut bacteria. *Microbiol Spectr* 5.
27. El Hage R, Hernandez-Sanabria E, Tom Van de Wiele (2017) Emerging trends in “smart probiotics”: Functional consideration for the development of novel health and industrial applications. *Front Microbiol* 8: 1889.
28. Soucy SM, Huang J, Gogarten JP (2015) Horizontal gene transfer: Building the web of life. *Nat Rev Genet* 16: 472-482.
29. Ochman H, Lawrence JG, Groisman EA (2000) Lateral gene transfer and the nature of bacterial innovation. *Nature* 405: 299-304.
30. Olivera PH, Touchon M, Cury J, et al. (2017) The chromosomal organization of horizontal gene transfer in bacteria. *Nat Comm* 8: 841.
31. Gilbert JA, Blaser MJ, Caporaso JG, et al. (2018) Current understanding of the human microbiome. *Nat Med* 24: 392-400.
32. Hayes CS, Aoki SK, Low DA (2010) Bacterial contact-dependent delivery systems. *Annu Rev Genet* 44: 71-90.
33. Manrique P, Dills M, Young MJ (2017) The human gut phage community and its implications for health and disease. *Viruses* 9: 141.
34. Acid tolerance of bacteria used in Doctors Biome™ DCUSA. Inc. subsidiary of Dupont, Danisco, USA.
35. Su GL, Ko CW, Bercik P, et al. (2020) AGA clinical practice guidelines on the role of probiotics in the management of gastrointestinal disorders. *Gastroenterology*.

DOI: 10.36959/621/560