

focus Mold detox like never before



Exceptionally pure, vigorous mycotoxin binder

Featuring:

- G-PUR[®] A highly purified natural zeolite. G-PUR[®] uses the clinoptilolite form, characterized by a higher silica level for maximum binding of mycotoxins, aluminum and ammonia.
- **PureBind™ Humic Acid** Exceedingly pure 70% humic acid (vs. standard 30%), featuring increased binding without common humic acid impurities.



Webinar Mold/Mycotoxin Exposure – the prevalence & suggested protocols

Watch on <u>https://youtu.be/</u> <u>MXF6TLFXH9w</u>

Mycotoxin detox protocol

Step 1 _ Prepare & open detox pathways

Tri-Fortify® Liposomal Glutathione -

Peer-reviewed, published, clinical research* confirms:

- 28% increase in red blood cell glutathione levels
- 400% increase in natural killer cell function
- 25% reduction in lipid peroxidation



Step 2 _ Bind & expel

MycoPul® – Advanced mycotoxin binder promotes healthy removal of multiple mycotoxins, as well as fungus, metals, and ammonia.

Transfer Factor Enviro™ – Promotes a healthy immune response to mold.

Core Minerals™ – 12 minerals + vitamin D3 to support essential minerals lost during detox.



- For all product sheets, see p.3 p.7
- Global Multi-Systemic treatment plan in mold & mycotoxins You can find the protocol at the end of this document

* Sinha R, Sinha I, Calcagnotto A, et al. Oral supplementation with liposomal glutathione elevates body stores of glutathione and markers of immune function. Eur J Clin Nutr. 2018;72(1):105-111. doi:10.1038/ejcn.2017.132

Tri-Fortify Watermelon[®] or Orange[®]



Detoxification with glutathione in high bioavailable formulation, powerful antioxidant, Natural Killer Cell support	
1 teaspoon (1 pack) per day, away from food	
236 ml per tube or 20 packs per box	
Glutathione Liposomal Vitamin C	450 mg 50 mg
	powerful antioxidant, Natural Killer Cell su 1 teaspoon (1 pack) per day, away from for 236 ml per tube or 20 packs per box Glutathione Liposomal







Oxidative stress markers Oxidized / Reduced GSH



Lipid Peroxidation (Reduced Cellular Membrane Oxidation)



Published research : Sinha, R., Sinha, I., Calcagnotto, A., Trushin, N. Oral supplementation with liposomal glutathione elevates body stores of glutathione and markers of immune function. Eur J Clin Nutr. 2018 Jan;72(1):105–111.



MycoPul™

indication	MycoPul is a combination of several binders, targeted to eliminate mycotoxins (toxins produced by molds), heavy metals and environmental toxins. MycoPul includes a patented, purified and natural form of zeolite.		
dosage	1 caps per day 1 hour before or 2 hours after eating		
packaging	30 caps per container		
composition (amount per 1 cap)	Activated Charcoal	150 mg	
	Fulvic acid	100 mg	
	Black humate powder	100 mg	
	Zeolite Clinoptilolite (G-Pur®)	100 mg	
	Microchitosan	100 mg	
	Bamboo extract	50 mg	

l information is exclusively aimed at and released to an audience of health care professiona



MycoPul[™] References

- ¹ El Khoury R, Choque E, El Khoury A, et al. Ota prevention and detoxification by actinobacterial strains and activated carbon fibers: Preliminary results. Toxins (Basel). 2018;10(4):1–16. doi:10.3390/toxins10040137
- ² Smyrna HP. G-PUR / Purified Clinoptilolite.; 2017.
- ³ Pavelić SK, Medica JS, Gumbarević D, Filošević A, Pržulj N, Pavelić K. Critical review on zeolite clinoptilolite safety and medical applications in vivo. Front Pharmacol. 2018;9(NOV):1–15.
- ⁴ Zhang L, Zeng Y, Cheng Z. Removal of heavy metal ions using chitosan and modified chitosan: A review. J Mol Liq. 2016;214:175–191.
- ⁵ Min LL, Zhong L Bin, Zheng YM, Liu Q, Yuan ZH, Yang LM. Functionalized chitosan electrospun nanofiber for effective removal of trace arsenate from water. Sci Rep. 2016;6(April).
- ⁶ Zia Q, Tabassum M, Gong H, Li J. A Review on Chitosan for the Removal of Heavy Metals lons. J Fiber Bioeng Informatics. 2019;12(3):103–128.
- ⁷ Jacqmin-Gadda H, Commenges D, Letenneur L, Dartigues JF. Silica and aluminum in drinking water and cognitive impairment in the elderly. Epidemiology. 1996;7(3):281–285.
- ⁸ Rondeau V, Jacqmin-Gadda H, Commenges D, Helmer C, Dartigues JF. Aluminum and silica in drinking water and the risk of Alzheimer's disease or cognitive decline: Findings from 15-year follow-up of the PAQUID cohort. Am J Epidemiol. 2009;169(4):489–496.
- ⁹ Davenward S, Bentham P, Wright J, et al. Silicon-rich mineral water as a non-invasive test of the "aluminum hypothesis" in Alzheimer's disease. J Alzheimer's Dis. 2013;33(2):423-430.
- ¹⁰ Dalton AJ. Commentary on "Non-invasive therapy to reduce the body burden of aluminium in Alzheimer's disease." J Alzheimer's Dis. 2006;10(1):29-31.
- Jones K, Linhart C, Hawkins C, Exley C. Urinary Excretion of Aluminium and Silicon in Secondary Progressive Multiple Sclerosis. EBioMedicine. 2017;26:60–67.



Transfer Factor Enviro[™]

indication	Specific Transfer Factors Targets specific pathogens
	Immune support = Multimessenger + Specific Transfer Factors
dosage	1 cap per day during the first week then take 2 caps per day away from food
packaging	60 gelcaps per container
composition (amount per 2 gelcaps)	Inositol (Rice bran) 261 mg Beta Glucan (Baker's yeast) Colostrum (Transfer Factors - pure amino acid sequences)



Core Minerals[™]

indication	Balanced mineral complex without iron: restorage of minerals during chelation therapy	
dosage	1 x 4 caps per day, 12 hours after chelation	
packaging	120 vegecaps per container	
composition	Calcium (as Calcium citrate + malate)	300 mg
(amount per 4 vegecaps)	Magnesium (as Magnesium glycinate)	300 mg
	Potassium (as Potassium citrate)	99 mg
	Zinc (as Zinc picolinate)	15 mg
	Copper (as Copper glycinate)	2 mg
	Manganese (as Manganese glycinate)	2 mg
	Boron (as Boron picolinate)	2 mg
	Chromium (as Chromium polynicotinate)	200 µg
	lodine	150 µg
	Selenium (as Selenium citrate)	100 µg
	Molybdenum (as Molybdenum picolinate)	100 µg
	Vanadium (as Vanadylsufate)	100 µg

Please find our referenced version on the professional section of our website. All information is exclusively aimed at and released to an audience of health care professionals.

Mold and Mycotoxins

Fungi are a large group of pathogens with two subgroups: yeast are single cell organisms, molds aremulticellular organisms. Molds reproduce by releasing spores. Some spores produce toxic mycotoxins.

Mold often grows in water damaged buildings. First step in treatment is removal from exposure (STEP 1). To remove the **mycotoxins** from the body, binders are used.

The toxic effects of mycotoxins include gastro-intestinal toxicity, inflammation, neuroinflammationand disruption of the immune response. Every treatment is an individualized interpretation on specific symptoms.

STEP 1 Removal from exposure	
STEP 2	Guttae Pepsini
Treatment of impaired intestinal barried induced by mycotoxins: Gut protocol	30 ml Dose: 3 x 10 - 20 drops at the start of the meal and with a small amount of water (swallow immediately)
	Gluten DPP IV Complex
	90 vcaps
	Dose: 3 x 1 caps per day, at the beginning of the meal
	Perm Plus Coated tablets
	90 coated tablets
	Dose: first month: 3 x 2 tablets per day then: 3 x 1 tablet per day, 20 minutes before the meal
	CoreBiotic
	<i>The use of soil-based probiotics to remote microbiome and reduce post-prandial raise in endotoxins</i> 60 vcaps
	Dose: 1 x 2 caps per day, at least 30 minutes before the meal, for minimum 2 months
STEP 3	Broccoraphanin Nrf2 Support
Detoxification support &	100 vcaps
rebuilding Glutathione levels	Dose: 1 caps per day
	Tri-Fortify Watermelon or Orange 236 ml
	Dose: 1 teaspoon/day, separated from meals



STEP 4

Immune support & reduce inflammation

Increasing NK Cell activity & support to the Regulatory T Cells

Multimessenger 90 caps Dose: 1 x 3 caps per day, just before breakfast

Specific activity targeting antigens

Transfer Factor Enviro 60 caps Dose: 2 caps before sleep

CytoQuel

90 vcaps Dose: 3 x 1 caps per day during or after meals

STEP 5

Reducing mycotoxin load in the body with binders.

Binders are not systemically absorbed.

Constipation occurs but binders rarely cause more severe symptoms. In a general way Binders should be taken separated from food, nutrition and drugs. Reducing mycotoxin load in the body with binders... etc. Combine binders based on general recommendations – see table

Carbon Black 400 mg (activated charcoal – certified grade)

200 vcaps Dose: 1-5 caps per day

Cholestyramine CSM Pure without additives 200 g powder

Dose: 2 to 4 x per day, max 4 g, dissolved in water or juice 1 hour before or 2 hours after meal or medication Go slow with the use of binders in hypersensitive patients Start by 1 x 1 gram – 1 x 2 grams, slowly increase when tolerated

Chlorella Glass grown 250 mg

200 vcaps Dose: Start slowly with 1 cap per day and gradually build up the daily dose 45-60 minutes prior to food

Mycopul

30 caps

Dose: 1 cap per day 1 hour before or 2 hours after meal

PectaSol-C

454 g powder Dose: 1 - 3 scoops per day in water or juice





References

Intestinal consequences

König, Julia, et al. "Randomized clinical trial: Effective gluten degradation by Aspergillus niger-derived enzyme in a complex meal setting." Scientific reports 7.1 (2017): 13100.

Luo, Su, et al. "In vitro and in vivo effects of a mycotoxin, deoxynivalenol, and a trace metal, cadmium, alone or in a mixture on the intestinal barrier." Environment international 132 (2019): 105082.

McFarlin, Brian K., et al. "Oral spore-based probiotic supplementation was associated with reduced incidence of post- prandial dietary endotoxin, triglycerides, and disease risk biomarkers." World journal of gastrointestinal pathophysiology 8.3 (2017): 117.

Ren, Zhihua, et al. "Progress in Mycotoxins Affecting Intestinal Mucosal Barrier Function." International journal of molecular sciences 20.11 (2019): 2777.

Thrasher JD, Crawley S. The biocontaminants and complexity of damp indoor spaces: more than what meets the eyes. Toxicol Ind Health. 2009 Oct-Nov.



Detoxification

C. A. Sun, L. Y. Wang, C. J. Chen et al., "Genetic polymorphisms of glutathione S-transferases M1 and T1 associated with susceptibility to aflatoxin-related hepatocarcinogenesis among chronic hepatitis B carriers: a nested case-control study in Taiwan," Carcinogenesis, vol. 22, no. 8, pp. 1289–1294, 2001

L. Alpsoy, A. Yildirim, and G. Agar, "The antioxidant effects of vitamin A, C, and e on aflatoxin B1-inducedoxidative stress in human lymphocytes," Toxicology and Industrial Health, vol. 25, no. 2, pp. 121–127, 2009

Morimitsu, Y., Nakagawa, Y. A Sulforaphane Analogue That Potently Activates the Nrf2-dependent Detoxification Pathway. J Biol Chem. 2002 Feb 1;277(5):3456-63.

Sinha, R., Sinha, I., Calcagnotto, A., Trushin, N. Oral supplementation with liposomal glutathione elevates body stores of glutathione and markers of immune function. Eur J Clin Nutr. 2018 Jan;72(1):105–111.

Woo, Kyung Jin, and Taeg Kyu Kwon. "Sulforaphane suppresses lipopolysaccharide-induced cyclooxygenase-2 (COX-2) expression through the modulation of multiple targets in COX-2 gene promoter." International immunopharmacology 7.13 (2007): 1776–1783.

Immune support

Birrell, Mark A., et al. "Resveratrol, an extract of red wine, inhibits lipopolysaccharide induced airway neutrophilia and inflammatory mediators through an NF-kB-independent mechanism." The FASEB journal 19.7 (2005): 840-841.

Chowdhury, Rupak, et al. "Curcumin attenuation of lipopolysaccharide induced cardiac hypertrophy in rodents." ISRN inflammation 2013 (2013).

Cognitive consequences

Anyanwu, Ebere C., Andrew W. Campbell, and Aristo Vojdani. "Neurophysiological effects of chronic indoor environmental toxic mold exposure on children." The Scientific World Journal 3 (2003): 281–290.

Casas, Lidia, et al. "Early life exposures to home dampness, pet ownership and farm animal contact and neuropsychological development in 4 year old children: A prospective birth cohort study." International journal of hygiene and environmental health 216.6 (2013): 690–697.

Centers for Disease Control and Prevention. "CDC estimates 1 in 59 children has been identified with autism spectrum disorder." (2018).

De Santis, Barbara, et al. "Role of mycotoxins in the pathobiology of autism: A first evidence." Nutritional neuroscience 22.2 (2019): 132-144.

Geschwind, Daniel H., and Matthew W. State. "Gene hunting in autism spectrum disorder: on the path to precision medicine." The Lancet Neurology 14.11 (2015): 1109–1120.

Gordon, Wayne A., et al. "Cognitive impairment associated with toxigenic fungal exposure: a replication and extension of previous findings." Applied Neuropsychology 11.2 (2004): 65–74.

Jedrychowski, Wieslaw, et al. "Cognitive function of 6-year old children exposed to mold-contaminated homes in early postnatal period. Prospective birth cohort study in Poland." Physiology & behavior 104.5 (2011): 989–995.

McCall, Robert B. "Childhood IQ's as predictors of adult educational and occupational status." Science 197.4302 (1977): 482-483.

Von Tobel, Jenny Sandström, et al. "Repeated exposure to Ochratoxin A generates a neuroinflammatory response, characterized by neurodegenerative M1 microglial phenotype." Neurotoxicology 44 (2014): 61-70.





Neuroinflammation

Liew, Winnie-Pui-Pui, and Sabran Mohd-Redzwan. "Mycotoxin: its impact on gut health and microbiota." Frontiers in cellular and infection microbiology 8 (2018): 60.

Ratnaseelan, Aarane M., Irene Tsilioni, and Theoharis C. Theoharides. "Effects of mycotoxins on neuropsychiatric symptoms and immune processes." Clinical therapeutics 40.6 (2018): 903–917.

Uetsuka, Koji. "Mechanisms of mycotoxin-induced neurotoxicity through oxidative stress-associated pathways." International journal of molecular sciences 12.8 (2011): 5213-5237.

Von Tobel, Jenny Sandström, et al. "Repeated exposure to Ochratoxin A generates a neuroinflammatory response, characterized by neurodegenerative M1 microglial phenotype." Neurotoxicology 44 (2014): 61-70.

Inflammation and immunity, oxidative stress

Al-Anati L, Petzinger E. Immunotoxicactivity of ochratoxin A. Journal of Veterinary Pharmacology and Therapeutics. 2006;29(2):79-90.

Campbell AW, Thrasher JD, Gray MR, Vojdani A. Mold and Mycotoxins: Effects on the Neurological and Immune Systems in Humans. Advances in Applied Microbiology. 2004:375–406.

Ermert D, Ram S, Laabei M. The hijackers guide to escaping complement: Lessons learned from pathogens. Molecular Immunology. 2019;114:49–61.

Hueza I, Raspantini P, Raspantini L, Latorre A, Górniak S. Zearalenone, an Estrogenic Mycotoxin, Is an Immunotoxic Compound. Toxins. 2014;6(3):1080–1095.

Hymery N, Sibiril Y, Parent-Massin D. In vitro effects of trichothecenes on human dendritic cells. Toxicology in Vitro. 2006;20(6):899-909.

Jahreis S, Kuhn S, Madaj A-M, Bauer M, Polte T. Mold metabolites drive rheumatoid arthritis in mice via promotion of IFN- gamma-and IL-17-producing T cells. Food and Chemical Toxicology. 2017;109:405-413.

Kankkunen P, Rintahaka J, Aalto A, et al. Trichothecene Mycotoxins Activate Inflammatory Response in Human Macrophages. The Journal of Immunology. 2009;182(10):6418-6425.

L. Alpsoy, A. Yildirim, and G. Agar, "The antioxidant effects of vitamin A, C, and e on aflatoxin B1-induced oxidative stress in human lymphocytes," Toxicology and Industrial Health, vol. 25, no. 2, pp. 121–127, 2009

Lehrnbecher T, Schmidt S. Why are natural killer cells important for defense against Aspergillus? Medical Mycology. 2019;57(Sup2).

Liew W-P-P, Mohd-Redzwan S. Mycotoxin: Its Impact on Gut Health and Microbiota. Frontiers in Cellular and Infection Microbiology. 2018;8.

Pestka JJ, Zhou H-R, Moon Y, Chung Y. Cellular and molecular mechanisms for immune modulation by deoxynivalenol and other trichothecenes: unraveling a paradox. Toxicology Letters. 2004;153(1):61-73.

Schütze N, Lehmann I, Bönisch U, Simon JC, Polte T. Exposure to Mycotoxins Increases the Allergic Immune Response in a Murine Asthma Model. American Journal of Respiratory and Critical Care Medicine. 2010;181(11):1188-1199.

Sherrington SL, Kumwenda P, Kousser C, Hall RA. Host Sensing by Pathogenic Fungi. Advances in Applied Microbiology. 2018:159-221.

Smith M, Mcginnis MR. Mycotoxins and their effects on humans. Clinical Mycology. 2009:649-656.



Vogl G, Lesiak I, Jensen D, et al. Immune evasion by acquisition of complement inhibitors: The mould Aspergillus binds both factor H and C4b binding protein. Molecular Immunology. 2008;45(5):1485–1493.

Wang H, Yadav JS. DNA damage, redox changes, and associated stress-inducible signalling events underlying the apoptosis and cytotoxicity in murine alveolar macrophage cell line MH-S by methanol-extracted Stachybotryschartarumtoxins. Toxicology and Applied Pharmacology. 2006;214(3):297-308.

Binders

Abbès S, et al. Preventive role of phyllosilicate clay on the Immunological and Biochemical toxicity of zearalenone in Balb/c mice. Int Immunopharmacol. 2006 Aug;6(8):1251-8.

Boylan, James J., John L. Egle, and Philip S. Guzelian. "Cholestyramine: use as a new therapeutic approach for chlordecone (Kepone) poisoning." Science 199.4331 (1978): 893–895.

Carfagna, Simona, et al. "Physiological and morphological responses of Lead or Cadmium exposed Chlorella sorokiniana 211-8K (Chlorophyceae)." SpringerPlus 2.1 (2013): 1-7.

Carson MS, Smith TK. (1983). Role of bentonite in prevention of T-2 toxicosis in rats. J Anim Sci, 57:1498–506

Cohn, William J., et al. "Treatment of chlordecone (Kepone) toxicity with cholestyramine: results of a controlled clinical trial." New England Journal of Medicine 298.5 (1978): 243–248.

D. E. Diaz, W. M. Hagler, J. T. Blackwelder et al., "Aflatoxin Binders II: reduction of aflatoxin M1 in milk by sequestering agents of cows consuming aflatoxin in feed," Mycopathologia, vol. 157, no. 2, pp. 233–241, 2004

Dvorak M. Ability of bentonite and natural zeolite to adsorb aflatoxin from liquid media. Vet Med (Praha), 1989. 34:307–16

Eliaz, Isaac, and Avraham Raz. "Pleiotropic Effects of Modified Citrus Pectin." Nutrients 11.11 (2019): 2619.

F. Galvano, A. Pietri, T. Bertuzzi, A. Piva, L. Chies, and M. Galvano, "Activated carbons: in vitro affinity for Ochratoxin A and deoxynivalenol and relation of adsorption ability to physicochemical parameters," Journal of Food Protection, vol. 61, no. 4, pp. 469–475, 1998

G. Avantaggiato, R. Havenaar, and A. Visconti, "Evaluation of the intestinal absorption of deoxynivalenol and nivalenol by an in vitro gastrointestinal model, and the binding efficacy of activated carbon and other adsorbent materials," Food and Chemical Toxicology, vol. 42, no. 5, pp. 817–824, 2004

Humphries, P., E. Pretorius, and H. Naude. "Direct and indirect cellular effects of aspartame on the brain." European journal of clinical nutrition 62.4 (2008): 451-462.

Miranda, M. S., Sunao Sato, and Jorge Mancini-Filho. "Antioxidant activity of the microalga Chlorella vulgaris cultered on special conditions." Bollettino chimico farmaceutico 140.3 (2001): 165–168.

Nakano, Shiro, Hideo Takekoshi, and Masuo Nakano. "Chlorella (Chlorella pyrenoidosa) supplementation decreases dioxin and increases immunoglobulin a concentration in breast milk." Journal of medicinal food 10.1 (2007): 134-142.

P. Wang, E. Afriyie-Gyawu, Y. Tang et al., "NovaSil clay intervention in Ghanaians at high risk for aflatoxicosis: II. Reduction in biomarkers of aflatoxin exposure in blood and urine," Food Additives and Contaminants Part A, vol. 25, no. 5, pp. 622–634, 2008.

Rafati-Rahimzadeh, Mehrdad, et al. "Current approaches of the management of mercury poisoning: need of the hour." DARU Journal of Pharmaceutical Sciences 22.1 (2014): 46.

Rafati-Rahimzadeh, Mehrdad, et al. "Current approaches of the management of mercury poisoning: need of the hour." DARU Journal of Pharmaceutical Sciences 22.1 (2014): 46.

Rycerz, Karol, and Jadwiga Elżbieta Jaworska-Adamu. "Effects of aspartame metabolites on astrocytes and neurons." Folia Neuropathol 51.1 (2013): 10-7.





Scaldaferri, Franco, et al. "Use and indications of cholestyramine and bile acid sequestrants." Internal and emergency medicine 8.3 (2013): 205-210.

Sikiru, Akeem Babatunde, et al. "Chlorella vulgaris supplementation effects on performances, oxidative stress and antioxidant genes expression in liver and ovaries of New Zealand White rabbits." Heliyon 5.9 (2019): e02470.

Tuchweber, Abdelhamid Kerkadi Claude Barriault Beatriz, and Andrzej A. Frohlich Ronald R. Marquardt. "Dietary cholestyramine reduces ochratoxin A-induced nephrotoxicity in the rat by decreasing plasma levels and enhancing fecal excretion of the toxin." Journal of Toxicology and Environmental Health Part A 53.3 (1998): 231–250.

Tuomi T, et al. Mycotoxins in crude building materials from water-damaged buildings. Appl Enviro Microbiology. 2000 May 1;66(5):1899-904.

